

Physics 566: Quantum Optics I

Problem Set #3: Solutions

Problem 1:

(a) Consider a statistical mixture

$$\hat{\rho} = P_+ | \uparrow_z \rangle \langle \uparrow_z | + P_- | \downarrow_z \rangle \langle \downarrow_z |$$

$$\text{where } P_{\pm} = \frac{1}{2} \left(1 \pm \frac{1}{\sqrt{2}} \right)$$

Density matrix $\rho_{ij} = \langle i | \hat{\rho} | j \rangle$

$$\text{In basis } |\pm_z\rangle \quad \hat{\rho} = \frac{1}{2} \begin{pmatrix} 1 + \frac{1}{\sqrt{2}} & 0 \\ 0 & 1 - \frac{1}{\sqrt{2}} \end{pmatrix}$$

$$\text{In basis } (| \uparrow_x \rangle = \frac{1}{\sqrt{2}} (| \uparrow_z \rangle + | \downarrow_z \rangle))$$

$$\hat{\rho} = \frac{1}{2} \begin{pmatrix} \langle \uparrow_x | \hat{\rho} | \uparrow_x \rangle & \langle \uparrow_x | \hat{\rho} | \downarrow_x \rangle \\ \langle \downarrow_x | \hat{\rho} | \uparrow_x \rangle & \langle \downarrow_x | \hat{\rho} | \downarrow_x \rangle \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 1 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & 1 \end{pmatrix}$$

Note: In this basis the density operator has off-diagonal elements. Nonetheless, it is a mixed state:

$$\text{Tr}(\hat{\rho}^2) = \frac{3}{4}$$

The Bloch vector can be seen immediately from the form in the z-basis.

$$\text{Tr}(\hat{\rho} \hat{\sigma}_x) = \text{Tr}(\hat{\rho} \hat{\sigma}_y) = 0$$

$$\text{Tr}(\hat{\rho} \hat{\sigma}_z) = P_{\uparrow_z} - P_{\downarrow_z} = \frac{1}{\sqrt{2}}$$

$$\Rightarrow \boxed{\vec{Q} = \frac{1}{\sqrt{2}} \vec{e}_z} \quad \text{Mixed state } |\vec{Q}| < 1$$

(b) Now we have a state

$$\hat{\rho} = \frac{1}{2} |\uparrow_{n_1}\rangle \langle \uparrow_{n_1}| + \frac{1}{2} |\uparrow_{n_2}\rangle \langle \uparrow_{n_2}|$$

$$\text{where } |\uparrow_n\rangle \langle \uparrow_n| = \frac{1}{2} (\hat{I} + \vec{e}_n \cdot \vec{\sigma}) \text{ from Prob 1}$$

$$\vec{e}_{n_1} = \frac{1}{\sqrt{2}} (\vec{e}_z \pm \vec{e}_x)$$

$$\Rightarrow \hat{\rho} = \frac{1}{2} \hat{I} + \frac{1}{4} (\vec{e}_{n_1} + \vec{e}_{n_2}) \cdot \vec{\sigma}$$

$$= \frac{1}{2} \hat{I} + \frac{1}{4} \left(\frac{2}{\sqrt{2}} \vec{e}_z \right) \cdot \vec{\sigma}$$

$$= \frac{1}{2} \left(\hat{I} + \frac{1}{\sqrt{2}} \vec{e}_z \right) \cdot \vec{\sigma} = \begin{bmatrix} \frac{1}{2}(1 + \frac{1}{\sqrt{2}}) & 0 \\ 0 & \frac{1}{2}(1 - \frac{1}{\sqrt{2}}) \end{bmatrix}$$

Same as $\hat{\rho}$ in part (b)!

Moral of the story: The ensemble decomposition is not unique. In fact, we can take any density matrix for a two-level system, described uniquely in terms of its Bloch vector \vec{Q} and

decompose it in terms of an ensemble of any two pure states described by unit vector \vec{e}_n with probability p_n if $\vec{Q} = p_{n_1} \vec{e}_{n_1} + p_{n_2} \vec{e}_{n_2}$.

(c) Two statistical mixtures

$$\hat{\rho}_1 = \sum_n p_n |\psi_n\rangle \langle \psi_n|$$

$$\hat{\rho}_2 = \sum_m q_m |\phi_m\rangle \langle \phi_m|$$

Aside: $|\psi_n\rangle \langle \psi_n| = \frac{1}{2} (\hat{I} + \hat{\sigma}_n)$ (where $\hat{\sigma}_n = \vec{e}_n \cdot \vec{\sigma}$)
Projector

$$\Rightarrow \hat{\rho}_1 = \underbrace{\left(\sum_n p_n \right)}_{= 1} \frac{1}{2} \hat{I} + \frac{1}{2} \left(\sum_n p_n e_n^2 \right) \cdot \vec{\sigma}$$

$$\Rightarrow \hat{\rho}_1 = \frac{1}{2} \hat{I} + \frac{1}{2} \vec{Q}_1 \cdot \vec{\sigma}$$

Similarly $\hat{\rho}_2 = \frac{1}{2} \hat{I} + \frac{1}{2} \vec{Q}_2 \cdot \vec{\sigma}$

where $\vec{Q}_2 = \sum_m q_m \vec{e}_m$

thus $\hat{\rho}_1 = \hat{\rho}_2 \quad \Leftrightarrow \quad \vec{Q}_1 = \vec{Q}_2$

Problem 2: Ambiguity of ensemble decomposition

Let $\hat{\rho}_1 = \sum_i p_i |\psi_i\rangle\langle\psi_i|$, $\hat{\rho}_2 = \sum_j q_j |\phi_j\rangle\langle\phi_j|$

Proof

$$\hat{\rho}_1 = \hat{\rho}_2 \text{ iff } \sqrt{q_j} |\phi_j\rangle = \sum_i y_{ji} \sqrt{p_i} |\psi_i\rangle$$

where y_{ji} are elements of unitary matrix.

Proof:

For convenience, define $|\bar{\Phi}_j\rangle = \sqrt{q_j} |\phi_j\rangle$

$$|\bar{\Psi}_i\rangle = \sqrt{p_i} |\psi_i\rangle$$

$$\therefore \langle \bar{\Phi}_j | \bar{\Phi}_j \rangle = q_j \quad \langle \bar{\Psi}_i | \bar{\Psi}_i \rangle = p_i$$

(1) Assume $|\bar{\Phi}_j\rangle = \sum_i y_{ji} |\bar{\Psi}_i\rangle$ y_{ji} elements of unitary matrix

Consider $\hat{\rho}_2 = \sum_j |\bar{\Phi}_j\rangle\langle\bar{\Phi}_j| = \sum_{ijk} y_{jk}^* y_{ji} |\bar{\Psi}_i\rangle\langle\bar{\Psi}_k|$

Aside: $(y_k)^* = U_{kj}^+$

$$\Rightarrow \hat{\rho}_2 = \sum_{ik} \underbrace{\left(\sum_j U_{kj}^+ y_{ji} \right)}_{\delta_{ik}} |\bar{\Psi}_i\rangle\langle\bar{\Psi}_k|$$

$$\Rightarrow \hat{\rho}_2 = \sum_i |\bar{\Psi}_i\rangle\langle\bar{\Psi}_i| = \hat{\rho}_1 \quad \checkmark$$

(ii) Now assume $\hat{P}_1 = \hat{P}_2 = \hat{P}$

\hat{P} being a Hermitian operator can be diagonalized

$$\Rightarrow \hat{P} = \sum_{\alpha} \lambda_{\alpha} |e_{\alpha}\rangle \langle e_{\alpha}|$$

where $\left\{ \begin{array}{l} \sum_{\alpha} \lambda_{\alpha} = 1 \text{ with } \lambda_{\alpha} \text{ real}, \quad 0 < \lambda_{\alpha} \leq 1 \\ \langle e_{\alpha} | e_{\beta} \rangle = \delta_{\alpha\beta} \end{array} \right.$

$$\text{let } |\bar{e}_{\alpha}\rangle = \sqrt{\lambda_{\alpha}} |e_{\alpha}\rangle \Rightarrow \hat{P} = \sum_{\alpha} |\bar{e}_{\alpha}\rangle \langle \bar{e}_{\alpha}|$$

$$\Rightarrow \sum_{\alpha} |\bar{\Psi}_{\alpha}\rangle \langle \bar{\Psi}_{\alpha}| = \sum_{\alpha} |\bar{e}_{\alpha}\rangle \langle \bar{e}_{\alpha}| = \sum_j |\bar{\Phi}_j\rangle \langle \bar{\Phi}_j|$$

We seek the relationship between

$$\{|\bar{\Psi}_{\alpha}\rangle\} \text{ and } \{|\bar{\Phi}_j\rangle\}$$

First note $\{|\bar{e}_{\alpha}\rangle\}$ form a basis for
the Hilbert space (with $\lambda_{\alpha}=0$ for
those vectors not in \hat{P})

$$\begin{aligned} \Rightarrow |\bar{\Psi}_{\alpha}\rangle &= \sum_{\alpha} |\bar{e}_{\alpha}\rangle \langle \bar{e}_{\alpha}| \bar{\Psi}_{\alpha}\rangle = \sum_{\alpha} |\bar{e}_{\alpha}\rangle \cdot \frac{\langle \bar{e}_{\alpha} | \bar{\Psi}_{\alpha}\rangle}{\sqrt{\lambda_{\alpha}}} \\ &= \sum_{\alpha} M_{\alpha} |\bar{e}_{\alpha}\rangle \end{aligned}$$

where $M_{\alpha} = \frac{\langle \bar{e}_{\alpha} | \bar{\Psi}_{\alpha}\rangle}{\sqrt{\lambda_{\alpha}}}$

$$\begin{aligned}
 \text{Now: } \sum_i M_{\alpha\beta} M_{i\beta}^* &= \sum_i \frac{\langle e_\alpha | \bar{\psi}_i \rangle \langle \bar{\psi}_i | e_\beta \rangle}{\sqrt{\lambda_\alpha \lambda_\beta}} \\
 &= \frac{1}{\sqrt{\lambda_\alpha \lambda_\beta}} \langle e_\alpha | \left(\sum_i |\bar{\psi}_i\rangle \langle \bar{\psi}_i| \right) |e_\beta \rangle \\
 &= \frac{1}{\sqrt{\lambda_\alpha \lambda_\beta}} \langle e_\alpha | \hat{p}_{e_\beta} \rangle = \frac{\lambda_\alpha S_{\alpha\beta}}{\sqrt{\lambda_\alpha \lambda_\beta}} = S_{\alpha\beta}
 \end{aligned}$$

\Rightarrow When arranged in a matrix, the columns of $M_{\alpha\beta}$ are orthonormal

(Subtle point: $M_{\alpha\beta}$ need not be square here, since # of pure states in the $\{|\bar{\psi}_i\rangle\}$ need not be the dimension of Hilbert space. However, we can always append extra columns in the orthogonal space to make $M_{\alpha\beta}$ unitary. Formally the matrix M is a "partial isometry".)

Thus since $|\bar{\psi}_i\rangle = \sum_\alpha M_{\alpha i} |e_\alpha\rangle$

$$|\bar{\psi}_j\rangle = \sum_\beta N_{j\beta} |\bar{e}_\beta\rangle$$

$$|\bar{\psi}_j\rangle = \sum_i U_{ji} |\bar{\psi}_i\rangle$$

where $U = NM^\dagger$ q.e.d.

Problem 3

The two dimensional vector space the specifies the polarization state of a photon defines a qubit. We make the association:

$$\vec{e}_+ = \frac{\vec{e}_H + i\vec{e}_V}{\sqrt{2}} \Rightarrow |\uparrow_z\rangle$$

$$\vec{e}_- = \frac{\vec{e}_H - i\vec{e}_V}{\sqrt{2}} \Rightarrow |\downarrow_z\rangle$$

$$(a) |\uparrow_x\rangle = \frac{|\uparrow_z\rangle + |\downarrow_z\rangle}{\sqrt{2}} \Leftrightarrow \frac{\vec{e}_+ + \vec{e}_-}{\sqrt{2}} \Rightarrow$$

$|\uparrow_x\rangle \Leftrightarrow \vec{e}_H \text{ (linear horizontal)}$
 $|\downarrow_x\rangle \Leftrightarrow i\vec{e}_V \equiv \vec{e}_V \text{ (linear vertical)}$


$$|\uparrow_y\rangle = \frac{|\uparrow_z\rangle + i|\downarrow_z\rangle}{\sqrt{2}} \Leftrightarrow \frac{\vec{e}_+ + i\vec{e}_-}{\sqrt{2}} \Rightarrow$$

$|\uparrow_y\rangle \Leftrightarrow \frac{1+i}{\sqrt{2}} \left(\frac{\vec{e}_H + \vec{e}_V}{\sqrt{2}} \right) = \frac{\vec{e}_H + \vec{e}_V}{\sqrt{2}} \text{ (linear at } 45^\circ \text{ between)} \quad (\vec{e}_H \text{ and } \vec{e}_V)$
 $|\downarrow_y\rangle \Leftrightarrow \frac{1-i}{\sqrt{2}} \left(\frac{\vec{e}_H - \vec{e}_V}{\sqrt{2}} \right) = \frac{\vec{e}_H - \vec{e}_V}{\sqrt{2}} \text{ (linear at } -45^\circ \text{ between)} \quad (\vec{e}_H \text{ and } \vec{e}_V)$

(b) For an arbitrary state of the qubit, $|\uparrow_n\rangle = \cos \frac{\theta}{2} |\uparrow_z\rangle + e^{i\phi} \sin \frac{\theta}{2} |\downarrow_z\rangle$, where (θ, ϕ) is the direction on the Poincaré sphere.

$$\Rightarrow |\uparrow_n\rangle = \cos \frac{\theta}{2} \vec{e}_+ + e^{i\phi} \sin \frac{\theta}{2} \vec{e}_-$$

Recall (e.g. see Jackson 3rd edition Chap 7.2), The polarization is generally elliptical



$$r \equiv \frac{\alpha_+}{\alpha_-} = \frac{\cos \frac{\theta}{2}}{\sin \frac{\theta}{2}} e^{i\phi} = \cot \frac{\theta}{2} e^{i\phi}$$

$$\text{Ratio of semimajor/semiminor axis} = \frac{1+r}{1-r} = \frac{1+\cot \frac{\theta}{2}}{1-\cot \frac{\theta}{2}} = \frac{1+\sin \theta}{1-\sin \theta}$$

The ellipticity is characterized by $||\alpha_+|^2 - |\alpha_-|^2| = |\cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2}| = |\cos \theta|$

The orientation of the ellipse is shown, making an angle $\phi/2$ w.r.t. \vec{e}_H .

(c) Sketch of the Poincaré sphere




Figure 2.7. Poincaré sphere.

(d) Wave plate induces a phase shift that differs for "ordinary" and "extra ordinary" polarization. The ordinary and extra ordinary directions are the eigenvectors. Thus, defining on the Poincaré sphere the vectors in Hilbert space

$$\vec{e}_o \Rightarrow | \uparrow_n \rangle, \quad \vec{e}_e \Rightarrow | \downarrow_n \rangle,$$

The waveplate performs the transformation

$$\hat{U}^{WP} = e^{i\phi_o} |\uparrow_n\rangle\langle\uparrow_n| + e^{i\phi_e} |\downarrow_n\rangle\langle\downarrow_n|$$

Note: An arbitrary $SU(2)$, $\hat{U} = e^{-i\frac{\phi}{2}} |\uparrow_n\rangle\langle\uparrow_n| + e^{i\frac{\phi}{2}} |\downarrow_n\rangle\langle\downarrow_n|$, for eigenvectors $|\uparrow_n\rangle$ & $|\downarrow_n\rangle$

We can factor out an overall phase that is irrelevant, $\frac{\phi_e + \phi_o}{2}$

$$\Rightarrow \hat{U}^{WP} = \cancel{e^{\frac{i(\phi_e + \phi_o)}{2}}} \left[e^{-i\frac{\Delta\phi}{2}} |\uparrow_n\rangle\langle\uparrow_n| + e^{+i\frac{\Delta\phi}{2}} |\downarrow_n\rangle\langle\downarrow_n| \right]$$

$$= e^{-i\frac{\Delta\phi}{2}} \hat{U}_n \quad \text{where } \Delta\phi = \phi_e - \phi_o$$

Now:

$$\vec{e}_o = \cos\theta \vec{e}_H + \sin\theta \vec{e}_V, \quad \vec{e}_e = -\sin\theta \vec{e}_H + \cos\theta \vec{e}_V \quad (\text{rotation on Poincaré sphere by } 2\theta \text{ around 3-axis}).$$

$$\text{With } \vec{e}_H \Rightarrow |\uparrow_x\rangle = |\uparrow_1\rangle, \quad \vec{e}_V = |\uparrow_x\rangle = |\downarrow_1\rangle$$

$$\Rightarrow |\uparrow_n\rangle = \hat{D}_3(2\theta) |\uparrow_1\rangle, \quad |\downarrow_n\rangle = \hat{D}_3(2\theta) |\downarrow_1\rangle$$

$$\Rightarrow \hat{U}_{\theta}^{\text{WP}}(\Delta\phi) = \hat{D}_3(2\theta) \hat{D}_1(\Delta\phi) \hat{D}_3^{+}(2\theta)$$

As a matrix, in the basis of $|\uparrow_x\rangle, |\downarrow_x\rangle$

$$\hat{D}_1(\Delta\phi) = e^{-i\frac{\Delta\phi}{2}\hat{\sigma}_1} \stackrel{=}{=} \begin{bmatrix} e^{-i\frac{\Delta\phi}{2}} & 0 \\ 0 & e^{+i\frac{\Delta\phi}{2}} \end{bmatrix}$$

$$\hat{D}_3(2\theta) = \cos\theta \hat{1} - i\sin\theta \hat{\sigma}_3 \stackrel{=}{=} \cos\theta \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - i\sin\theta \underbrace{\begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}}_{\hat{U}_3 \text{ in } \{|\uparrow_1\rangle, |\downarrow_1\rangle\}} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$$

$$\Rightarrow \hat{U}_{\theta}^{\text{WP}}(\Delta\phi) \stackrel{=}{=} \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} e^{-i\frac{\Delta\phi}{2}} & 0 \\ 0 & e^{+i\frac{\Delta\phi}{2}} \end{bmatrix} \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$$

$$\Rightarrow \hat{U}_{\theta}^{\text{WP}}(\Delta\phi) \stackrel{=}{=} \begin{bmatrix} \cos\left(\frac{\Delta\phi}{2}\right) + i\cos 2\theta \sin\left(\frac{\Delta\phi}{2}\right) & -i\sin\left(\frac{\Delta\phi}{2}\right) \sin 2\theta \\ -i\sin\left(\frac{\Delta\phi}{2}\right) \sin 2\theta & \cos\left(\frac{\Delta\phi}{2}\right) - i\cos 2\theta \sin\left(\frac{\Delta\phi}{2}\right) \end{bmatrix}$$

This is a familiar transformation from optics, which specifies the input-output relation $\vec{E}_{\text{out}} = \alpha_{\text{in}} \vec{e}_H + \beta_{\text{in}} \vec{e}_V \Rightarrow \vec{E}_{\text{out}} = \alpha_{\text{out}} \vec{e}_H + \beta_{\text{out}} \vec{e}_V$

$$\begin{bmatrix} \alpha_{\text{out}} \\ \beta_{\text{out}} \end{bmatrix} = \hat{U}_{\theta}^{\text{WP}}(\Delta\phi) \begin{bmatrix} \alpha_{\text{in}} \\ \beta_{\text{in}} \end{bmatrix}$$

(c) A quarter-wave plate, $L = \frac{\lambda}{4(n_e - n_o)}$ $\Rightarrow \Delta\phi = \frac{\pi}{2}$

$$\Rightarrow U_{QWP} = \begin{bmatrix} \frac{1+i\cos 2\theta}{\sqrt{2}} & \frac{-i}{\sqrt{2}} \sin 2\theta \\ \frac{-i}{\sqrt{2}} \sin 2\theta & \frac{1-i\cos 2\theta}{\sqrt{2}} \end{bmatrix}$$

To transform linear and \vec{e}_H to \vec{e}_V : $\begin{bmatrix} 1 \\ 0 \end{bmatrix} \Rightarrow \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ i \end{bmatrix}$
 $\begin{bmatrix} 0 \\ 1 \end{bmatrix} \Rightarrow \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -i \end{bmatrix} = \frac{-i}{\sqrt{2}} \begin{bmatrix} i \\ 1 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} i \\ 1 \end{bmatrix}$

\Rightarrow Choose $\Theta = \frac{\pi}{4}$ (45° between fast & slow axes) \Rightarrow


$U_{QWP}(\Theta = \frac{\pi}{4}) = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & i \\ i & 1 \end{bmatrix}$ in $\vec{e}_H \vec{e}_V$ basis \Rightarrow Rotation by $-\frac{\pi}{2}$ around 2-axis of Poincaré

A half-wave plate, $L = \frac{\lambda}{2(n_e - n_o)}$ $\Rightarrow \Delta\phi = \pi$


$$\Rightarrow U_{QWP} = -i \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{bmatrix}$$

To transform $\vec{e}_H \rightarrow \vec{e}_V$: $\begin{bmatrix} 1 \\ 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ We achieve this

by choosing $\Theta = \frac{\pi}{4}$. The transformation is a rotation about 2-axis by π .



Quarter-wave plate



Half-Wave plate

From this geometric construction we see immediately how to orient the wave plate. The axis of rotation (the eigenvector of the waveplate) should be half way between the $\vec{e}_H + \vec{e}_V$ directions. The quarter wave plate then is a $\frac{\pi}{2}$ rotation on the Poincaré sphere, corresponding to $\vec{e}_H \Rightarrow \vec{e}_+^+$, $\vec{e}_V \Rightarrow \vec{e}_-^-$. The half-wave plate maps $\vec{e}_H \Rightarrow \vec{e}_V$, $\vec{e}_V \Rightarrow \vec{e}_H$ (overall phase irrelevant).

(f) From part (d), the SU(2) rotation corresponding to a wave plate, with the crystal axes oriented at angle θ w.r.t \vec{e}_H, \vec{e}_V direction is $U_{\theta}^{WP}(\Delta\phi) = e^{-i\frac{\Delta\phi}{2}\vec{e}(\theta)\cdot\hat{\sigma}} = \cos\frac{\Delta\phi}{2}\hat{1} - i[\cos 2\theta \hat{\sigma}_1 + \sin 2\theta \hat{\sigma}_2] \sin\frac{\Delta\phi}{2}$. (Note: A subtle point — in part (d) we wrote the matrices in the \vec{e}_H, \vec{e}_V basis this defines \hat{G}_1 in the usual Poincaré sphere) Thus, for a quarter waveplate and half waveplate respectively:

$$\text{QWP: } U_{\theta}^{WP}(\frac{\pi}{2}), \quad \text{HWP: } U_{\theta}^{WP}(\pi)$$

We seek to show that we can construct an arbitrary SU(2) rotation on the Poincaré sphere with two QWP and one HWP. To do this I will employ an Euler angle parameterization. Recall

$$U \in \text{SU}(2) \Rightarrow \exists \alpha, \beta, \gamma \text{ (Euler angles) st. } U = \hat{D}_3(\alpha) \hat{D}_2(\beta) \hat{D}_3(\gamma) \quad \begin{pmatrix} \text{rotation about 3-axis, then} \\ \text{2-axis, then 3-axis} \end{pmatrix}$$

(Note this is one Euler decomposition)

Thus a sequence of QWP-HWP-QWP

$$\begin{aligned} & \Rightarrow U_{\theta_a}^{WP}(\frac{\pi}{2}) U_{\theta_b}^{WP}(\pi) U_{\theta_c}^{WP}(\frac{\pi}{2}) \\ &= [\hat{D}_3(2\theta_a) \hat{D}_1(\frac{\pi}{2}) \hat{D}_3^+(2\theta_a)] [\hat{D}_3(2\theta_b) \hat{D}_1(\pi) \hat{D}_3^+(2\theta_b)] [\hat{D}_3(2\theta_c) \hat{D}_1(\frac{\pi}{2}) \hat{D}_3^+(2\theta_c)] \\ &= \hat{D}_3(2\theta_a) \hat{D}_1(\frac{\pi}{2}) \hat{D}_3(2\theta_b - 2\theta_a) \underbrace{\hat{D}_1(\pi) \hat{D}_3(2\theta_c - 2\theta_b)}_{\text{flip sign with } \pi\text{-rotation}} \hat{D}_1(\frac{\pi}{2}) \hat{D}_3^+(2\theta_c) \\ &= \hat{D}_3(2\theta_a) \hat{D}_1(\frac{\pi}{2}) \hat{D}_3(2\theta_b - 2\theta_a) \underbrace{\hat{D}_3(-2\theta_c + 2\theta_b)}_{\text{flip sign with } \pi\text{-rotation}} \hat{D}_1(-\pi) \hat{D}_1(\frac{\pi}{2}) \hat{D}_3^+(2\theta_c) \\ &= \hat{D}_3(2\theta_a) \hat{D}_1(\frac{\pi}{2}) \hat{D}_3(2\theta_b - 2\theta_a) \hat{D}_3(-2\theta_c + 2\theta_b) \hat{D}_1^+(\frac{\pi}{2}) \hat{D}_3^+(2\theta_c) \\ &= \hat{D}_3(2\theta_a) \underbrace{\hat{D}_1(\frac{\pi}{2}) \hat{D}_3(4\theta_b - 2\theta_a - 2\theta_c)}_{\text{rotate 2} \rightarrow 1} \hat{D}_1^+(\frac{\pi}{2}) \hat{D}_3^+(2\theta_c) \\ &= \hat{D}_3(2\theta_a) \underbrace{\hat{D}_2(4\theta_b - 2\theta_a - 2\theta_c)}_{\alpha} \underbrace{\hat{D}_3(-2\theta_c)}_{\gamma} \quad \text{Q.E.D. Phew!} \end{aligned}$$