

# Lecture 6


## (Reflection, Interference and Standing Waves)

Physics 262-01 Spring 2019

Douglas Fields

# Un-bounded Periodic Wave Description

- We can put these together:



$$y(x,t) = A \cos(kx - \omega t)$$


$$k = \frac{2\pi}{\lambda}, \quad \omega = \frac{2\pi}{T}$$

Wave travelling to the right

# Reading Quiz

- At what value of  $x(>0)$  do all of these waves have the same phase?

- A)  $x=3$
- B)  $x=6.2$
- C)  $x=9.5$
- D)  $x=12.5$



# Energy and Power

- I want to cover a little of energy transfer in a wave, although I find the discussion in the book mostly boring.
- Please don't memorize the equation in the book, but let's do go through the derivation.
- The work done on a piece of a string by a wave is just as usual:  $dW = \vec{F} \cdot d\vec{y}$


where, the force  $F$  is the total force on the piece, and  $dy$  is the little displacement it undergoes.

# Power transmitted by a mechanical wave

- As before, but now just examining what happens at one point,

For small slopes,  $T_x \approx T$ , so

$$T_{Ly} = -T \frac{\partial y}{\partial x}$$



- So, the work is just:

$$dW = -T \frac{\partial y}{\partial x} dy$$

- And the power (work per time) is:

$$P(x, t) = \frac{dW}{dt} = -T \frac{\partial y(x, t)}{\partial x} \frac{\partial y(x, t)}{\partial t}$$

# Power transmitted by a mechanical wave

$$P(x,t) = \frac{dW}{dt} = -T \frac{\partial y(x,t)}{\partial x} \frac{\partial y(x,t)}{\partial t}$$

- But,

$$y(x,t) = A \cos(kx - \omega t) \Rightarrow$$

$$\frac{\partial y(x,t)}{\partial x} = -Ak \sin(kx - \omega t)$$

$$\frac{\partial y(x,t)}{\partial t} = A\omega \sin(kx - \omega t)$$

- So,

$$P(x,t) = TA^2 k \omega \sin^2(kx - \omega t)$$
$$= \sqrt{\mu T} A^2 \omega^2 \sin^2(kx - \omega t)$$

$$v = \frac{\omega}{k} = \sqrt{\frac{T}{\mu}} \Rightarrow$$

$$\frac{k}{\omega} = \sqrt{\frac{\mu}{T}}$$

# Average Power

$$P(x, t) = \sqrt{\mu T} A^2 \omega^2 \sin^2(kx - \omega t)$$


- And the average of  $\sin^2$  is just  $1/2$ , so:

$$P_{avg} = \frac{1}{2} \sqrt{\mu T} A^2 \omega^2$$


- What to get from this?
- For mechanical waves, the power is a function of both the amplitude and the frequency squared.
- For E&M waves, it is independent of the frequency (as we will discover).

# Waves at Boundaries

- What happens to a wave when it reaches a boundary depends upon the boundary:
  - Fixed boundary (no energy transfer):



- Unfixed boundary (no energy transfer)





# Waves at Boundaries

- What happens when there *can* be energy transfer at a boundary?
  - Wave from low mass/length to high mass/length:



- Wave from high mass/length to low mass/length:



- Remember by taking the limit as heavier string becomes a wall...

# Waves at Boundaries

- What happens when there the wave is bounded on two sides?



# Superposition

- The wave equation, because it is linear, lends itself to superposition of solutions:

- If  $y_1$  is a solution of the wave equation:

$$y_1(x, t) = \cos(k_1x - \omega_1t)$$


- And  $y_2$  is also a solution of the wave equation:

$$y_2(x, t) = \cos(k_2x - \omega_2t)$$

- Then  $(Ay_1 + By_2)$  is also a solution of the wave equation.

# Superposition

- Superposition also means that waves “interfere” with each other.
  - If two different waves both exist on a string (or any other medium), then the resultant wave is just the superposition (linear sum) of the individual waves.
  - This can mean the amplitude gets bigger,
  - Or smaller...
- But each wave exists on the string independently!



# Standing Waves

- Let's take  $y_1$  as a wave moving to the right:

$$y_1(x, t) = A \cos(kx - \omega t)$$

- And  $y_2$  as a wave moving to the left:

$$y_2(x, t) = A \cos(kx + \omega t)$$

- Then  $(y_1 + y_2)$  is also a solution of the wave equation:

$$y_1(x, t) + y_2(x, t) = A \cos(kx - \omega t) + A \cos(kx + \omega t)$$

# Standing Waves

- Lets simplify this:

$$y_1(x, t) + y_2(x, t) = A \cos(kx - \omega t) + A \cos(kx + \omega t)$$

$$\text{But, } \cos(A) + \cos(B) = 2 \sin \frac{1}{2}[A + B] \cos \frac{1}{2}[A - B]$$

So,

$$\begin{aligned} y_1(x, t) + y_2(x, t) &= 2A \sin \frac{1}{2}[(kx - \omega t) + (kx + \omega t)] \cos \frac{1}{2}[(kx - \omega t) - (kx + \omega t)] \\ &= 2A \sin(kx) \cos(\omega t) \end{aligned}$$

# Standing Waves


- Examine this solution:

$$\begin{aligned}y_1(x, t) + y_2(x, t) &= 2A \sin(kx) \cos(\omega t) \\ &= \underbrace{[2A \cos(\omega t)]}_{\text{time dep. amplitude}} \underbrace{\sin(kx)}_{\text{position dep.}}\end{aligned}$$

- In position, there is a sin dependence whose amplitude depends upon time.


# Standing Waves

$$\begin{aligned}
 y_1(x,t) + y_2(x,t) &= 2A \sin(kx) \cos(\omega t) \\
 &= \underbrace{[2A \cos(\omega t)]}_{\text{time dep. amplitude}} \underbrace{\sin(kx)}_{\text{position dep.}}
 \end{aligned}$$



$$y_1(x,t)$$

$$y_2(x,t)$$



$$y_1(x,t)$$

$$y_2(x,t)$$

